



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

368. Proposed by G. I. HOPKINS, Professor of Mathematics and Astronomy, Manchester High School, Manchester, N. H.

It is required to construct the triangle having given, base, vertical angle, and ratio of its altitude to sum of the other two sides.

369. Proposed by W. J. GREENSTREET, A. M., Editor, *Mathematical Gazette*, Stroud, England.

Prove by inversion that if two circles cut at a given angle, touch each a given circle, and pass each through the same fixed point, then shall the envelope of the points of contact be a conic.

370. Proposed by R. C. ARCHIBALD, Paris, France.

The trisectors of the angles of any triangle ABC are, in order, AF , AE , CE , CD , BO , BF . Show synthetically that D , E , F are the vertices of an equilateral triangle.

CALCULUS.

295. Proposed by C. E. FLANNAGAN, Wheeling, W. Va.

A hawk can fly v feet per second, a hare can run v' feet per second. The hawk, when a feet vertically above the hare, gives chase and catches the hare when the hare has run b feet. Find the length of the curve of pursuit. [Echols' *Differential and Integral Calculus*, page 253, Ex. 20.]

296. Proposed by C. N. SCHMALL, New York City.

Two currents C_1 and C_2 produce deflections ϕ_1 , ϕ_2 , respectively, in a tangent galvanometer. When is $(\phi_1 - \phi_2)$ a maximum?

MECHANICS.

249. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

A load P is supported by three strings of equal size attached at the vertices of a triangle sides a , b , c lying in a horizontal plane. The load is vertically under the centroid of the triangle at a distance h from it. Find the stresses in the strings.

250. Proposed by C. N. SCHMALL, New York City.

A smooth circular table is surrounded by a smooth vertical rim. A ball of elasticity e is projected from a point at the rim in a line making an angle ϕ with the radius through that point. Show that the ball will return to the starting point after the second impact if

$$\tan \phi = \sqrt{\frac{e^3}{e^2 + e + 1}}.$$

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

174. Proposed by B. KRAMER, Student, University of Pittsburg, Pittsburg, Pa.

Find a general solution of $x(x+a)=y^2$, a , x , and y being integers. Given a , required to find x to satisfy conditions.